



Assessing the achievements and uncertain future of paleoepigenomics

Rick WA Smith¹  & Amy L Non^{*,2} 

¹Department of Sociology and Anthropology, George Mason University, Fairfax, VA 22030, USA

²Department of Anthropology, University of California, San Diego, La Jolla, CA 92093, USA

*Author for correspondence: alnon@ucsd.edu

“Although ancient epigenomic data can provide exciting hints of the past, this research is at its best when it integrates data from ancient and living human genomics/epigenomics, history and archaeology, which creates a more holistic view of human evolutionary history and our past environments.”

First draft submitted: 4 October 2021; Accepted for publication: 11 November 2021; Published online: 1 December 2021

Keywords: ancient DNA • ancient epigenomics • DNA methylation • paleoepigenomics

Paleoepigenomics is an emerging area of research that focuses on generating epigenomic data from ancient or extinct organisms with the goal of understanding their regulatory, environmental and evolutionary significance [1–3]. We are excited about the prospects of learning more about epigenetic regulation in the past, which may help us understand environmental plasticity, drivers of inter- and intraspecies variation, and the evolution of modern and ancient diseases, among other phenotypes. However, we are also concerned about the potential for speculation in an area of research that has often been overhyped. Given the complex combination of methods involved in this work as well as the significant gaps that remain in our understanding of regulatory mechanisms, there are reasons to maintain a healthy skepticism about what paleoepigenomics may achieve. The history of paleogenomics has been no stranger to the speculative and the far-fetched, with some early claims of DNA from the Mesozoic turning out to be contamination [4]. Similarly, epigenomics has sometimes been home to the sensational and the brash, perhaps most notably in the ongoing debates regarding transgenerational inheritance and the significant gaps that remain between identifying epigenomic changes and understanding their functional roles in health outcomes [5,6].

Paleogenomics and epigenomics are both still relatively new areas of science, and their respective histories of overreach and overhype have led to the development of rigorous methodological, authentication and analytical standards by which to guide and scrutinize research [4,7–10]. Paleoepigenomics is the integration of these two fields, which not only brings a double burden of credibility by existing standards but also raises new questions of plausibility. Understandably, there has been substantial enthusiasm around the potential of using paleoepigenomics to elucidate ancient environments, lifeways, and evolutionary events. However, we question whether the field is ready to dive into fully interpreting the epigenomes of the past when conventional epigenomics is still figuring out fundamental questions of the relevance and role of the epigenome for diseases and other phenotypes among living organisms in the present. Here we briefly assess paleoepigenomic work to date in humans and closely related hominins and consider the exciting advancements that have been made in the field. In addition, we consider the known limitations of paleoepigenomic research today and urge caution if – and until – the science catches up with the hype.

Paleoepigenomics: methods, applications & achievements

Early on, much of the work in paleoepigenomics was necessarily methodological and proof-of-concept based. Although epigenetic studies in extant organisms focus on a variety of epigenetic regulations at the DNA, RNA and protein levels, the majority of paleoepigenetic/omic research has focused on cytosine methylation. To our knowledge, the first evidence demonstrating that cytosine methylation status could be detected in ancient DNA (aDNA) came from the observation that enzymatic removal of deaminated cytosines depended on dinucleotide context [11].

DNA degrades rapidly after an organism dies and is characterized by highly fragmented nucleotide chains and spontaneous hydrolytic deamination of cytosines [1]. The presence of random postmortem transitions of cytosines into other pyrimidines (i.e., thymine or uracil) results in higher sequencing errors for aDNA relative to nondegraded sources of DNA. To increase sequencing accuracy, enzymatic repair techniques, such as uracil-DNA-glycosylase (UDG) treatment, have been used to remove cytosine-to-uracil transitions and have become common in aDNA library preparations. However, Briggs *et al.* found that some deaminated cytosines resist enzymatic removal and that this resistance depends strongly on dinucleotide context, with CpG dinucleotides retaining a higher fraction of deaminated cytosines after UDG treatment than other dinucleotide contexts (i.e., CpA, CpC and CpT) [11]. Because cytosines degrade differently after death, with unmethylated cytosines degrading to uracils and methylated cytosines degrading to thymines [12], the persistence of deaminated cytosines in CpG dinucleotides after repair reflects the presence of deaminated 5mCs (i.e., thymines) that are not removed by UDG treatment.

This finding led to the development of computational methods for deamination-dependent cytosine methylation prediction, which have become the most widely used techniques in paleoepigenomics. Here we focus our discussion on these deamination-dependent methods but note that some paleoepigenomic studies have used conventional bisulfite sequencing approaches as well [12,13]. The application of deamination-dependent approaches led to the first methylome reconstructions for an ancient human [14], Neandertals and Denisovans [15], and the inference of methylation in an increasing variety of other ancient and extinct organisms [14–16]. These methods have since been formalized into computational pipelines and open-source software programs for methylome prediction in aDNA [17]. In addition, methods for methylated binding domain enrichment and array-based methylation detection have been evaluated for use in aDNA [16,18].

One of the greatest promises of paleoepigenomics was that it might provide novel insights into major evolutionary changes and adaptive transitions in humans [2,3]. Some progress has been made toward these goals, with a few studies beginning to identify morphological and potentially behavioral differences between humans and our closest primate and hominin relatives. Following the sequencing of the complete Neandertal genome, studies began to identify coding and regulatory differences between living humans and Neandertals. One such difference is a polymorphism identified in the miRNA miR-1304, which is derived in living humans and is predicted to increase the number of putative regulatory targets by more than tenfold. Among the predicted regulatory targets are genes involved in neurodevelopment and enamel formation, and it has been suggested that this polymorphism may play a role in the evolution of dental and behavioral distinctions between living humans and Neandertals [19]. Building on the analysis of sequence-based regulatory differences, the reconstruction of Neandertal and Denisovan methylomes allowed for the identification of thousands of potentially differentially methylated regions between contemporary and archaic humans, including differences in *HOXD* cluster methylation, which might explain morphological differences such as limb proportions between living humans and Neandertals [3,15].

More recently, methylation maps have been used to predict the anatomy of Denisovans, which is currently known only from fragmentary and very limited subfossil remains. These analyses suggest that Denisovans likely shared features with Neandertals, such as robust jaws, long and low craniums, low foreheads, thick enamel, large rib cages, wide pelvises and large femoral articulations. However, these analyses also identified 11 features potentially distinct in Denisovans, including an elongated dental arch, changes in the dimensions of the mandible and lateral expansion of the parietal bones in the cranium. In the case of the Denisovan mandible, Gokhman *et al.* reported that seven out of eight of their *a priori* morphological predictions based on epigenetic differences matched the description of the first confirmed Denisovan mandible [20]. Subsequently, epigenetic changes have been linked with potential changes in the facial, vocal, spinal and mandibular anatomy of living humans since our geographic distancing from Neandertals and Denisovans [2,21].

Some of the first epigenetic studies of past human lifeways identified potential epigenetic differences related to diet and immune function between groups that vary in their subsistence patterns, including across the Mesolithic to Neolithic transition. For example, Gokhman *et al.* identified hypermethylation of *LOC654433*, *RBM46* and *EXD3* and hypomethylation of *BOLA3* in ancient hunter-gatherers compared with sedentary peoples [22]. More recently, Seguin-Orlando *et al.* detected some evidence of differential methylation in *CIQb* and *LCK* – genes that are related to humoral immune complement response and T-cell maturation, respectively – between Mesolithic and Neolithic peoples in France [23]. These types of studies can provide intriguing hints of past life experiences and their regulatory impacts but are also relevant for understanding variation in methylation related to diets and disease states among living people.

Taken together, work in paleoepigenomics has demonstrated not only that epigenetic marks can be reconstructed but also that they can lend some insights into ancient environments and evolutionary processes. The field has therefore accomplished some of its original goals. However, it is also important to be cognizant of the technical limitations of this work and consequently what questions are – and may remain – out of reach.

Limitations

The demonstrated correspondence between *a priori* morphological predictions of Denisovan anatomy and newly reported subfossil remains is one of the strongest pieces of evidence to date supporting the viability of paleoepigenomics for predicting unknown phenotypic outcomes [20]. However, it is important to keep in mind Gokhman *et al.*'s own note of caution that their study relied on predicting broad directional changes in anatomical features rather than precise phenotypic outcomes. In addition, although this represents an incredibly exciting result, the emergence of corroborating subfossil evidence represents a fortuitous and perhaps nonreplicable set of circumstances. How might we vet paleoepigenomic predictions in other contexts where there may be no possibility of corroborating evidence of phenotypes and no means to assess the regulatory pathways that help shape them? This would at a minimum require a more robust understanding of methylation and its specific regulatory consequences among extant peoples, ideally living in very similar conditions. Unfortunately, we have yet to fully arrive at such an understanding, and this places constraints on what can be inferred from methylation patterns in the past. In paleoepigenomics, we are often left with only the baseline epigenetic marks – with greatly diminished access to or knowledge of the downstream biology – and therefore little to no reliable way to mechanistically link epigenetic patterns to phenotypic outcomes.

The relationship between methylation and expression is remarkably complex. Even in studies of living organisms it is difficult to meaningfully link epigenetic marks with their regulatory impacts, and correlations can vary by genomic context, cell or tissue type [24]. Because of this, our ability to interpret the specific function of ancient methylation patterns is very limited. We should therefore be exceedingly cautious to not overinterpret results. This is true of any paleoepigenetic/omic study no matter the phenotype under study but is perhaps especially true in the case of predicting cognitive or behavioral distinctions between groups because the linkage between epigenetic marks and behavioral outcomes is especially tenuous.

In addition, constraints caused by gaps in the archaeological record and uneven DNA preservation in the available archaeological contexts place profound limits on both the types of questions that can be addressed and the sample sizes and statistical power available to analyze them. Although ancient methylation from a small number of Neandertals provides a unique glimpse into the past, these few individuals do not represent all the variation across an entire group, nor allow us to search for subtle trait- or disease-related methylation patterns in a way that satisfies the current analytical standards of the field. For example, statisticians have recently indicated that sample sizes upwards of 1000 people may be necessary to ensure adequate power to detect small disease-associated epigenetic differences when using array-based epigenome-wide approaches [8]. Further, fragmentary skeletal preservation, even in a rich archaeological context, may hinder interpretations of any detected methylation differences. If a past population experienced famine, violence and deprivation – all of which could potentially influence methylation in similar and overlapping ways – it can become impossible to disentangle effects. We face similar limitations in studies with living people even when we can directly measure these simultaneous exposures, but these problems are exacerbated in ancient contexts where exposures can only be inferred from limited remains.

Paleoepigenomic analyses are also limited to the tissues that preserve most often in the archaeological record – bone, teeth and hair. It is thus somewhat unsurprising when studies of ancient methylomes highlight skeletal differences of Neandertals and Denisovans or when methylome data from hair demonstrate patterns characteristic of that tissue [14,15,20]. Because the epigenome varies substantially across tissues, this raises the question of what insights can be achieved when we lack some of the most physiologically relevant tissues in the archaeological record. For example, if a goal is to understand a phenotype such as stress response in archaeological communities but the most commonly studied tissues of blood and buccal cells are not preserved, can reconstruction of methylation in less physiologically relevant tissues such as bone, teeth and hair suffice? These issues have been thoughtfully considered by Gokhman *et al.* [3] but continue to represent significant hurdles to the reconstruction of ancient environments and lifeways, a core goal of the field.

The ability of paleoepigenomics to address more specific questions requiring fine-scale precision will necessitate the development of methods for enhanced methylome resolution in aDNA. Although the deamination-dependent technique has become the most widely used method, it is very limited in resolution, typically providing regional

methylation estimates in genomic windows of a few hundred base pairs. This approach is therefore constrained to analyses of differentially methylated regions rather than individual CpGs and only in individuals and genomic regions where sufficient deamination has occurred to infer methylation patterns. Thus, a differentially methylated region approach may determine regional differences in methylation but is not well suited to finer-scale epigenomic questions, such as determining the methylation status of transcription factor binding sites in the promoter regions of dietary or stress-related genes, which sometimes entails determining the methylation state of single cytosines. Bisulfite pyrosequencing of aDNA is the most direct method for detecting ancient methylation patterns at single-cytosine resolution. However, this method is highly destructive to already heavily degraded aDNA and thus far has only been successfully applied to reconstruct methylation of repetitive elements rather than the single-copy loci that would be necessary for specific analyses of diet and stress, for example [12,13]. Importantly, even with this direct measurement method, postmortem deamination can still bias methylation estimates. Methylation levels may be artificially reduced or show increased variability depending on aDNA preservation and the amount of cytosine deamination in the genome [12].

Future potential

Keeping these limitations in mind, we recognize the potential value of paleoepigenomic techniques, particularly to answer questions that can only be addressed in ancient or archaeological contexts. Such questions could include some of the impacts of historical traumas, such as those resulting from slavery and forced removals of African and Indigenous peoples from their homelands. Events such as these may carry forward to impact the health of living descendants today via the perpetuation over generations of harmful social and environmental conditions that maintain epigenetic states, the potential (yet undemonstrated) mechanisms of transgenerational inheritance through the germline, or other mechanisms. To distinguish historic effects from those of more recent experiences, methylation levels assessed close to the time of the historic trauma would be necessary. As previously noted, however, this would require either remarkable aDNA preservation, such as in recent historical archaeological contexts, or the development of higher resolution methods for methylome reconstruction. In addition, although paleoepigenomic data may elucidate some of the molecular mechanisms associated with historical traumas, it is important for researchers working in this area to be cognizant that many impacts of historical traumas are already well-characterized by multiple lines of historical, biomedical and community knowledge, among others. Epigenomics researchers of the past, like those of the present, should also be careful not to introduce or reinforce new forms of biomedical reductionism by overemphasizing a deterministic role of the environment or 'victim blaming' of parents or marginalized groups. Given the combination of paleogenomic and epigenomic approaches involved in this work, future research should also be mindful of the necessity for ethical and decolonial approaches to archaeological and aDNA research with historically marginalized people [25,26] as well as critical scholarship considering the potential and pitfalls of epigenomics as a means to elucidate historical traumas [27].

Another area of future potential for paleoepigenomics is the opportunity within ancient contexts to detect effects of longer-term exposures than can be measured among living people, such as the stress of living through lengthy sociopolitical transformations, climatic disruptions, environmental degradations, or generations of exposure to lead poisoning or other toxins. It is also now possible to detect epigenomic effects of past exposures in living adults that may no longer be observable or measurable in other ways, such as smoke exposure *in utero*, remarkably, over 20 years prior [28], or even in a previous generation Luo *et al.* [29]. Perhaps paleoepigenomics may ultimately enable inference of other unmeasurable exposures, such as past extreme climates, with the potential to predict epigenomic consequences of current and future climate change. Finally, it is clear that paleoepigenomics may be able to provide additional morphological guidelines for identifying new subfossil evidence of species that are known only from highly fragmentary remains.

Recommendations

The future of paleoepigenomics will depend on advances in our knowledge of contemporary epigenomics, which can refine our approach on where to look and exactly what to look for in the past (i.e., tissues and regions of the genome most affected by particular exposures). Inferences of past cellular phenotypes could be improved once we have more extensive documentation of the relationship between DNA methylation and expression across genomic regions in contemporary studies and the role of other epigenetic mechanisms, such as histone modifications, miRNAs and availability of transcription factors, that also influence expression patterns. These advances can be

aided by the expansion of existing public databases that document epigenetic associations (EWAS Atlas [30]) and tissue-specific gene expression (Genotype-Tissue Expression project [31]).

In terms of study design and implementation, we recommend that paleoepigenomic reconstructions be considered only in circumstances where DNA preservation and genome coverage are exceptionally high. This constrains analyses to the occasional ancient individual, to a limited set of environmental contexts (e.g., cold, arid) that especially facilitate aDNA preservation, and to more recent historic archaeological contexts where adequate DNA preservation may be more common. Because of the high risk and ethical stakes of this research, we recommend that paleoepigenomic research be considered on the back end of other, lower-risk genomic work, such as analyses of population history or natural selection, if and when the DNA preservation allows. To preserve precious samples and ancestral remains, we further recommend ancient epigenetic/omic studies be limited to questions that cannot be answered by studies of living people. For example, just as with genomics research, many questions about our species' deep evolutionary history can be inferred using comparative approaches between living human and nonhuman primates [32] and by using complex computational modeling techniques.

Conclusion & future perspective

The history of early research in both paleogenomics and epigenomics has been fraught with speculative claims, with subsequent and ongoing developments of methodological, authentication, and analytical criteria. Merging these two often overhyped fields entails a double burden of credibility. We urge caution and suggest that researchers avoid conclusions that overreach their data. Finally, we emphasize that just as with paleogenomic data, there are limits to the insights that can be gained through paleoepigenomic data, as samples and ancestral remains are scarce and unevenly preserved, and evolutionary and historic events can never be directly observed. Although ancient epigenomic data can provide exciting hints of the past, this research is at its best when it integrates data from ancient and living human genomics/epigenomics, history and archaeology, which creates a more holistic view of human evolutionary history and our past environments.

Author contributions

RWA Smith and AL Non contributed equally to the writing and editing of this manuscript.

Acknowledgments

The authors thank the anonymous reviewers for their helpful feedback in revising the manuscript.

Disclosure

The funders played no role in the content of the manuscript.

Financial & competing interests disclosure

The authors acknowledge support from Vanderbilt University (Discovery Grant) and the Wenner-Gren Foundation (to RWA Smith). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as: ● of interest; ●● of considerable interest

1. Liu Y, Weyrich LS, Llamas B. More arrows in the ancient DNA quiver: use of paleoepigenomes and paleomicrobiomes to investigate animal adaptation to environment. *Mol. Biol. Evol.* 37(2), 307–319 (2020).
- **Suggests paths for integrating paleoepigenomic data with other metagenomic approaches.**
2. Mathov Y, Batyrev D, Meshorer E, Carmel L. Harnessing epigenetics to study human evolution. *Curr. Opin. Genet. Dev.* 62, 23–29 (2020).
3. Gokhman D, Meshorer E, Carmel L. Epigenetics: it's getting old. Past meets future in paleoepigenetics. *Trends Ecol. Evol.* 31(4), 290–300 (2016).
4. Pääbo S, Poinar H, Serre D *et al.* Genetic analyses from ancient DNA. *Annu. Rev. Genet.* 38, 645–679 (2004).
5. Non AL. Social epigenomics: are we at an impasse? *Epigenomics* 13(21), 1747–1759 (2021).
- **Provides a recent overview of progress and barriers in social epigenomic research, with recommendations for improving future studies in this field.**

6. Heard E, Martienssen RA. Transgenerational epigenetic inheritance: myths and mechanisms. *Cell* 157(1), 95–109 (2014).
7. Llamas B, Willerslev E, Orlando L. Human evolution: a tale from ancient genomes. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 372(1713), 20150484 (2017).
8. Mansell G, Gorrie-Stone TJ, Bao Y *et al.* Guidance for DNA methylation studies: statistical insights from the Illumina EPIC array. *BMC Genomics* 20(1), 366 (2019).
9. Salas LA, Peres LC, Thayer ZM *et al.* A transdisciplinary approach to understand the epigenetic basis of race/ethnicity health disparities. *Epigenomics* 13(21), 1761–1770 (2021).
10. Llamas B, Valverde G, Fehren-Schmitz L, Weyrich LS, Cooper A, Haak W. From the field to the laboratory: controlling DNA contamination in human ancient DNA research in the high-throughput sequencing era. *Sci. Technol. Archaeol. Res.* 3(1), 1–14 (2017).
11. Briggs AW, Stenzel U, Meyer M, Krause J, Kircher M, Paabo S. Removal of deaminated cytosines and detection of *in vivo* methylation in ancient DNA. *Nucleic Acids Res.* 38(6), e87 (2010).
- **Provided the first evidence that it was possible to detect DNA methylation status in aDNA based on cytosine deamination patterns.**
12. Smith RW, Monroe C, Bolnick DA. Detection of cytosine methylation in ancient DNA from five Native American populations using bisulfite sequencing. *PLoS One* 10(5), e0125344 (2015).
13. Llamas B, Holland ML, Chen K, Cropley JE, Cooper A, Suter CM. High-resolution analysis of cytosine methylation in ancient DNA. *PLoS One* 7(1), e30226 (2012).
14. Pedersen JS, Valen E, Velazquez AM *et al.* Genome-wide nucleosome map and cytosine methylation levels of an ancient human genome. *Genome Res.* 24(3), 454–466 (2014).
- **Landmark study that was the first to reconstruct an epigenome for an ancient human.**
15. Gokhman D, Lavi E, Prufer K *et al.* Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science* 344(6183), 523–527 (2014).
- **First study to reconstruct methylomes for closely related hominins and to compare differences in methylation between these hominins and living humans.**
16. Seguin-Orlando A, Gamba C, Sarkissian C *et al.* Pros and cons of methylation-based enrichment methods for ancient DNA. *Sci. Rep.* 5, 11826 (2015).
17. Hanghøj K, Renaud G, Albrechtsen A, Orlando L. DamMet: ancient methylome mapping accounting for errors, true variants, and post-mortem DNA damage. *Gigascience* 8(4), giz025 (2019).
- **Formalized a computational pipeline for inferring DNA methylation from cytosine deamination patterns in ancient genomes and developed an open-source software program for paleoepigenomic analysis.**
18. Schmidt M, Maixner F, Hotz G *et al.* DNA methylation profiling in mummified human remains from the eighteenth-century. *Sci. Rep.* 11(1), 15493 (2021).
19. Lopez-Valenzuela M, Ramírez O, Rosas A *et al.* An ancestral miR-1304 allele present in Neanderthals regulates genes involved in enamel formation and could explain dental differences with modern humans. *Mol. Biol. Evol.* 29(7), 1797–1806 (2012).
20. Gokhman D, Mishol N, De Manuel M *et al.* Reconstructing Denisovan anatomy using DNA methylation maps. *Cell* 179(1), 180–192.e10 (2019).
- **First study to estimate anatomy based on paleoepigenomic data and to successfully predict an unknown ancient phenotype.**
21. Batyrev D, Lapid E, Carmel L, Meshorer E. Predicted archaic 3D genome organization reveals genes related to head and spinal cord separating modern from archaic humans. *Cells* 9(1), 48 (2020).
22. Gokhman D, Malul A, Carmel L. Inferring past environments from ancient epigenomes. *Mol. Biol. Evol.* 34(10), 2429–2438 (2017).
- **First paleoepigenomic study to investigate epigenomic variation related to past lifeways in humans and to identify diet-related methylation differences between ancient hunter-gatherers and sedentary people.**
23. Seguin-Orlando A, Donat R, Der Sarkissian C *et al.* Heterogeneous hunter-gatherer and Steppe-related ancestries in late Neolithic and Bell Beaker genomes from present-day France. *Curr. Biol.* 31(5), 1072–1083.e10 (2021).
24. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.* 13(7), 484–492 (2012).
25. Wagner JK, Colwell C, Claw KG *et al.* Fostering responsible research on ancient DNA. *Am. J. Hum. Genet.* 107(2), 183–195 (2020).
26. Dunnivant J, Justinvil D, Colwell C. Craft an African American graves protection and repatriation act. *Nature* 593(7859), 337–340 (2021).
27. Kowal E, Warin M. Anthropology, indigeneity, and the epigenome. *Am. Anthropol.* 120(4), 822–825 (2018).
28. Richmond RC, Suderman M, Langdon R, Relton CL, Davey Smith G. DNA methylation as a marker for prenatal smoke exposure in adults. *Int. J. Epidemiol.* 47(4), 1120–1130 (2018).
29. Luo R, Zhang H, Mukherjee N *et al.* Association of grandmaternal smoking during pregnancy with DNA methylation of grandchildren: the Isle of Wight study. *Epigenomics* 13(18), 1473–1483 (2021).

30. Li M, Zou D, Li Z *et al.* EWAS Atlas: a curated knowledgebase of epigenome-wide association studies. *Nucleic Acids Res.* 47(D1), D983–D988 (2019).
31. Carithers LJ, Moore HM. The Genotype-Tissue Expression (GTEx) project. *Biopreserv. Biobank.* 13(5), 307–308 (2015).
32. Jeong H, Mendizabal I, Berto S *et al.* Evolution of DNA methylation in the human brain. *Nat. Commun.* 12(1), 2021 (2021).